

FE DE ERRATA

En el trabajo “Estudio experimental sobre el efecto regulador de Qinggan Dongyin en la homeostasis de los linfocitos T en ratones MRL/lpr”, publicado en el Vol. 66 número 3, septiembre de 2025, es necesario hacer unas correcciones a saber: en las páginas 274 y 275 deben sustituirse las Tablas 1 y 2 por esta nueva versión.

ERRATUM

In the article “Experimental study on the regulatory effect of Qinggan Dongyin on T lymphocyte homeostasis in MRL/lpr mice”, published in Vol. 66, Issue 3, September 2025, the following corrections are necessary: on pages 274 and 275, Tables 1 and 2 should be replaced with this updated version.

Table 1. Distribution of CD4 + T, CD8 + T, Th1, Th2, Th17 and Treg cells in the spleen of mice in each group.

	Control group (n=6)	Model group (n=6)	QGDY group (n=6)
CD4+ T (%)	19.67 ± 0.63	15.37 ± 0.21***	17.52 ± 0.32###
CD8+ T (%)	12.65 ± 0.55	6.38 ± 0.29***	9.49 ± 0.17###
Th1 (%)	0.82 ± 0.16	3.61 ± 0.39**	1.60 ± 0.06##
Th2 (%)	0.75 ± 0.07	3.38 ± 0.35***	1.58 ± 0.03###
Th17 (%)	1.36 ± 0.20	2.83 ± 0.10***	1.97 ± 0.13###
Treg (%)	2.59 ± 0.08	0.69 ± 0.30**	1.61 ± 0.04##

Note: Cells of CD4-positive T cells (CD4+ T), CD8-positive T cells (CD8+ T), T-helper 1 cells (Th1), T-helper 2 cells (Th2), T-helper 17 cells (Th17), Regulatory T cells (Treg) in model and treatment group compared with the control group using one-way ANOVA method, and the mean ± standard deviation (SD) is used for the result representation. **p<0.01, ***p<0.001; compared with the model group, ###p<0.001, ##p<0.01.

Table 2. Concentrations of IFN-γ, IL-6, TNF-α, IL-17A, TGF-β and PFP in the plasma of mice in each group.

	Control group (n=6)	Model group (n=6)	QGDY group (n=6)
IFN-γ (ng/L)	113.07 ± 31.21	457.42 ± 218.77*	166.81 ± 22.34 [#]
IL-6 (pg/mL)	28.07 ± 8.76	80.29 ± 39.76*	40.66 ± 47.38
TNF-α (ng/L)	80.11 ± 44.98	373.12 ± 146.68**	186.70 ± 111.87 [#]
IL-17A (pg/mL)	17.69 ± 7.18	108.23 ± 38.61**	50.77 ± 32.63 [#]
TGF-β (ng/L)	27.20 ± 18.38	177.47 ± 78.95**	69.97 ± 26.65 [#]
PFP (ng/L)	47.41 ± 14.01	14.64 ± 8.78**	34.95 ± 20.50

Note: The cytokines of Interferon-gamma (IFN-γ), Interleukin-6 (IL-6), Tumor Necrosis Factor-α (TNF-α), Interleukin-17A (IL-17A), Transforming Growth Factor-beta (TGF-β), Perforin (PFP) in model and treatment groups compared with the control group using One-way ANOVA method, and the mean ± standard deviation (SD) is used for the result representation. *p<0.05, **p<0.01; compared with the model group, [#]p<0.05.